



CULTIVATION OF SALT-TOLERANT CROPS IN SALINE SOIL CONDITIONS DUE TO GLOBAL CLIMATE CHANGE

B. N. Atamurodov 1,

O. M. Yusupova 2

1Senior Lecturer (PhD) of the Department of Irrigation and Land Reclamation,
Bukhara State Technical University, Uzbekistan

2Student of Bukhara State Technical University, Uzbekistan

Abstract

Crop is considered the most valuable, high – yield grain crop food, fodder, technical and agrotechnical importance. As food, cornstarch's Dogi is used. His Dogi is considered very nutritious and has an average content of 10,6% cutlet, 1,4% ash substances. But, in corn grain, the protein content is low. For this reason, bread is closed by adding 25-30% bug'doy flour to the corn flour. Since corn Dogi has a lot of precipitation (4,3-5,0%) in its composition, it quickly hurts it. Grain porridge is separated in special machines, flour is prepared from the rest, because in the composition of corn porridge contains about 25 – 40% fat, from which the oil used for food is prepared. In addition, cereals from corn cereals are prepared, during the milk – wax ripening period, it can be used as food, in which it is fried (tonsils) and boiled.

Keywords: Corn, high yield, fodder, nutritious, corn Dani, corn flour.

Introduction

Currently, attention is being paid to improving the land reclamation situation in the agricultural and water sector, increasing the efficiency of canals and collector-trenches in the supply of water to the irrigated areas of farmer farms. Shular in addition, the effective use of Land-Water Resources requires the improvement of modern resurstejamkor irrigation techniques and technologies. In the strategy of action on further development of the Republic of Uzbekistan "... in order to increase the competitiveness of the national economy, it is necessary to reduce the consumption of energy and resources, to introduce a wide range of resource-intensive technologies into production." In this regard, it is important to carry out



research work on increasing the fertility of lands on the basis of the development and application of resurstejamkor irrigation methods of crops and the water use plan. The world is engaged in purposeful research on the use of land and water resources and improvement of the melioration condition of the soil. In this regard, the implementation of large-scale anti-erosion research in irrigated areas, the creation of techniques and technologies for the protection of land and Water Resources, the introduction of soil-protecting crops into the structure of arable areas is one of the important tasks in the field of implementation with oats. The development of livestock is one of the most important issues in the provision of environmentally friendly food products to the population, increasing the stock of fodder for livestock.

Corn is a valuable, fertile, cereal crop in multi-band use. From cornstarch it is possible to get a good concentrate, Blue Mass and silage. 1 kg of grain contains 1,34 units of nutrients. Silage, prepared when cornstarchs enter the milk-wax ripening period, is the best food for livestock. At 100 kg of silage there are 24 feed units. In the preparation of land for planting corn, phosphorus (ammonium per 150-200 kg/ha or suprifos per 300-350 kg/ha) fertilizer is given using the technique of spraying NRU-0,5 rubles of fertilizer before planting. If there is a sufficient amount of rotten manure in the farmer's farm, it is ground with the help of fertilizing techniques. Driving on the ground is carried out at a depth of 25-28 CM. In cases where the surface of the field is uneven, it is leveled with special techniques. Work on eliminating unevenness in a small area can be carried out by hanging rollers on the tractor. After that, the field is made of molasses. Sowing corn on the grain in the spring (for corn, which will be 120 days, at this time, when sown, farmer farms achieve a high result from grain yield to 3-4 t/ha. The time of germination should not remain until may, otherwise the yield will decrease. The planting period is 10-15 April for the regions located in the Central Region (Tashkent, Sirdarya, Jizzakh, Samarkand and Fergana Valley regions), 2 weeks before that (25-30 march) in the southern regions, and 2 weeks after that (25-30 April) in the northern regions. In the spring term, agrotechnical activities in the planting and maintenance of corn for silage, even seedsarfi, are conducted as they are planted in the grain. Depending on the ripening period, the number of seedlings will vary (for goats – 50-55 thousand Bush/ha; for the middle-late-70 thousand Bush/ha; for the middle-fairy tale and Fairy Tales-80 thousand Bush / ha).



For maize silage, it is considered effective to plant in the spring from intermediate crops or from grain crops with a spike in the summer (fast-growing hybrids of maize at this time, and the highest grain mass is obtained. If the fast-growing hybrids of corn are sown in the spring term, then the yield will sharply decrease to 2-3 t/ha then it is possible to plant. It is recommended to plant corn only for silage in the northern regions of our republic on the fields that are free of Spike crops. In the spring planting of silos and grain is recommended late-maturing, late-maturing varieties and hybrids of corn ("Uzbekistan-601 ESV", "Uzbekistan-420 VL"), early-maturing, early-maturing varieties and hybrids for cereals in summer ("Karasuv-350 AMV", "Uzbekistan-300 MV"). When using pneumatic SPCH-6 seyralka, the planting depth is 5-6 cm. it consists of 22-25 kg seeds per hectare. Bunda the size of each row is 60 cm, 70 cm and 90 cm. Lee will. 60 cm. li is placed in the matriarchs between 21 cm, 24 cm and 30 cm, and in the matriarchs between 24 cm and 30 cm. 70 cm. li is placed in the matriarchs between 18 cm and 20 cm, and the matriarchs between 26 cm. 90 cm. li is placed in the matriarchs between 14 cm, 16 cm and 20 cm, and in the matriarchs between 16 cm and 20 cm. Seed germination is given if there is a lack of moisture in the ground, so that the seeds germinate after planting. In the spring, after the seeds are planted in the ground, there is a lot of precipitation, the areas on which the resin is formed are surface cultivated with the help of cultivators, or the resin is softened with special adaptations. If, after sowing the seeds, the humidity on the ground is high and the air temperature drops, the seeds will rot. In this case, re-planting is carried out.

When there are 3-4 leaves after planting, the first cultivar is done, manual chopsticks are done, if there is a large number of seedlings, the yagana work is carried out. When there are a lot of weeds, they are treated with special herbicides. When 5-6 leaves are harvested in corn, the first irrigation is carried out by laying 150-180 kg of urea or 250-300 kg of ammonium nitrate per hectare together with ration. When there are 8-10 leaves on the stem, the second cultivar is made. If the field is pressed by a weed, chupik works are carried out, when 10-12 leaves are formed, watered for the second time along with the second feeding. The fasting of corn is watered for the third time to keep the moisture in the norm during the period when it comes to flowering. This watering will last longer than other irrigation. If during this period there is no emphasis on the norm of irrigation, productivity (40-50%) will decrease sharply. This period interval is 7-10 days. At this time,



pollination is carried out on the corn crop. It has been proven based on scientific experiments that give well-pollinated plants a high level yield. In corn, the fourth watering is carried out when there is a period of ripening of milk-wax in the grain, at this time the grains are filled and contain the necessary nutrients. At this time, the highest mass and nutrient is achieved if it is rolled into silage. During the ripening of milk-wax in the silo mowing use Maral, KPI-2,4, e-280-duty combines. It is checked before planting corn on the grain, if the plant is in a period of mummification, it is allowed to harvest tavarbop cereals. Seeds grown on large fields (elite, o'l hybrids) are carried out when full ripening begins. In this case, Khersones–7, Khersones–9 are used. The work that is carried out before planting corn in the summer consists in the following: fertilizing the ground with phosphorus fertilizer (150-200 kg/ha ammofos or 300-350 kg/ha suprifos), plowing the land (at a depth of 25-28 CM), leveling, scratching and molting. In summer planting, it is recommended to have all the activities that are applied in the spring, and seedling the same. From the hybrids of the corn farmer in the morning ("Karasuv–350 AMV" and "Uzbekistan–300 MV"), it is recommended to plant until 20-25 June and to obtain silos until 15 July in order to obtain fully ripe grain. Cultivation of corn in angiosperms in the summer agrotechnical measures are carried out like spring planting. It should be taken into account that corn, which is planted in the summer, is rapidly growing. Due to this, it is recommended that the plant quickly undergoes processing, feeding and watering between rows.

In summary, it can be said that the development of water-intensive irrigation technology of prospective maize varieties in the conditions of alluvial soils of the meadow of Bukhara region is an urgent issue, taking into account the importance of growing maize and its use in various sectors of the public economy, in the field of pharmaceuticals, cattle breeding and others, scientific research "Uzbekistan-2001", Borja"of maize, which is planted as the main crop in the soil conditions, Scientific-research work on the cultivation of the"Pioneer" variety is carried out, preliminary results are achieved, and scientific research work in this direction is continuing.



REFERENCES

1. Atamurodov, B. N., Sobirov, K. S., & Najmiddinov, M. M. (2022). BASICS OF FARMING ON SALINE AND SALINE-PRONE SOILS. *Oriental renaissance: Innovative, educational, natural and social sciences*, 2(6), 725-730.
2. Xamidova, S. M., Juraev, U. A., & Atamurodov, B. N. (2022). EVALUATION OF THE EFFECTIVENES OF PHYTOMELIORATIVE MEASURES IN THE TREATMENT OF RECLAMATION OF SALINE SOILS. *Web of Scientist: International Scientific Research Journal*, 3(6), 835-841.
3. Jurayev, A. K., Jurayev, U. A., Atamurodov, B. N., Sobirov, K. S., & Najmiddinov, M. M. (2022). IRRIGATION OF COTTON BY WATER-SAVING METHOD. *Oriental renaissance: Innovative, educational, natural and social sciences*, 2(6), 718-724.
4. Atamurodov, B. N., Sobirov, K. S., & Najmiddinov, M. M. (2022). USE OF RESOURCE-EFFICIENT IRRIGATION TECHNOLOGY IN THE REPUBLIC OF UZBEKISTAN. *Science and innovation*, 1(D2), 96-100.
5. Jurayev, A. K., Jurayev, U. A., Atamurodov, B. N., Najmiddinov, M. M., & Sobirov, K. S. (2022). EFFECTIVE USE OF WATER IN IRRIGATED AREAS. *Oriental renaissance: Innovative, educational, natural and social sciences*, 2(6), 810-815.
6. Jurayev, A. K., Jurayev, U. A., Atamurodov, B. N., Sobirov, K. S., & Najmiddinov, M. M. (2022). GROWING TOMATOES HYDROPONICALLY IN GREENHOUSES. *Science and innovation*, 1(D2), 87-90.
7. Atamurodov, B. N., Murodov, O. U., Najmiddinov, M. M., & Sobirov, K. S. (2022). IN IRRIGATION OF AGRICULTURAL CROPS, IRRIGATION WITH DIFFERENT QUALITY WATER. *Science and innovation*, 1(D2), 91-95.
8. Jurayev, A. K., Jurayev, U. A., Atamurodov, B. N., Sobirov, K. S., & Najmiddinov, M. M. (2022). SOYBEANS ARE TRANSPLANTED INTO SALINE AND SALINE SOILS TO JUSTIFY THE EFFECTIVENESS OF DRIP IRRIGATION.
9. Jurayev, A. K., Jurayev, U. A., Atamurodov, B. N., Sobirov, K. S., & Najmiddinov, M. M. (2022). IRRIGATION OF GOOSE BY WATER-SAVING METHOD.



10. Jurayev, A. K., Jurayev, U. A., Atamurodov, B. N., Sobirov, K. S., & Najmiddinov, M. M. (2022). SCIENTIFIC AND PRACTICAL IMPORTANCE OF EFFICIENT USE OF WATER IN IRRIGATED LAND.
11. Jurayev, A. Q., Jurayev, U. A., Atamurodov, B. N., & Najmiddinov, M. M. (2021). Cultivation of Corn as a Repeated Crop. *European Journal of Life Safety and Stability* (2660-9630), 10, 49-51. Jurayev, A. Q.,
12. Jurayev, U. A., Atamurodov, B. N., & Najmiddinov, M. M. (2021). Scientific Benefits and Efficiency of Drip Irrigation. *Journal of Ethics and Diversity in International Communication*, 1(6), 62-64.
13. Jurayev, A. Q., Jurayev, U. A., Atamurodov, B. N., & Najmiddinov, M. M. (2021). Aphorisms of Farming in the Method of Kidroponics. *International Journal of Discoveries and Innovations in Applied Sciences*, 1(6), 133-135.
14. Jo'rayev, U. A., Jo'rayev, A. Q., & Atamurodov, B. N. (2021). Application of Provided Irrigation Technologies in Irrigated Agriculture. *International Journal of Development and Public Policy*, 1(6), 164-166.
15. Atamurodov, B. N., Ibodov, I. N., Najmiddinov, M. M., & Najimov, D. Q. The Effectiveness of Farming in the Method of Hydroponics. *International Journal of Human Computing Studies*, 3(4), 33-36.
16. Jurayev, A. Q., Jurayev, U. A., Atamurodov, B. N., & Najmiddinov, M. M. (2021). The Main Purpose of Drip Irrigation in Irrigation Farming and Its Propagation. *European Journal of Life Safety and Stability* (2660-9630), 10, 46-48.
17. Atamurodov, B. N., & Jo'rayev, U. A. (2024, October). QIYIIN MELIORATSIYALANADIGAN YERLARNING MELIORATIV HOLATINI YAXSHILASHDA INNOVATSION SHO'R YUVISH TEXNOLOGIYASINING AHAMIYATI. In *Uz-conferences* (No. 1, pp. 133-138).
18. Juraev, A. K., Khamidov, M. K., Juraev, U. A., Atamurodov, B. N., Murodov, O. U., Rustamova, K. B., & Najmiddinov, M. M. (2024). The effect of deep softeners on irrigation, salt washing and cotton yield on soils with heavy mechanical composition. In *BIO Web of Conferences* (Vol. 103, p. 00017). EDP Sciences.
19. Jurayev, A. K., Jurayev, U. A., Atamurodov, B. N., Sobirov, K. S., & Najmiddinov, M. M. (2022). WATERING THEIR CROPS WITH WATER OF



DIFFERENT QUALITY. *Oriental renaissance: Innovative, educational, natural and social sciences*, 2(6), 1251-1257.

20. Atamurodov, B. N., Sobirov, K. S., & Najmiddinov, M. M. (2022). Rational Use of Water in Agricultural Regions. *Miasto Przyszłości*, 25, 88-89.

21. Khamidov, M., Juraev, A., Juraev, U., Atamurodov, B., Rustamova, K., Najmiddinov, A., & Nurbekov, A. (2022, July). Effects of deep softener and chemical compounds on mechanical compositions in heavy, difficult-to-ameliorate soils. In *IOP Conference Series: Earth and Environmental Science* (Vol. 1068, No. 1, p. 012017). IOP Publishing.

22. Jurayev, A. K., Jurayev, U. A., Atamurodov, B. N., Sobirov, K. S., & Najmiddinov, M. M. (2022). WATERING THEIR CROPS WITH WATER OF DIFFERENT QUALITY. *Oriental renaissance: Innovative, educational, natural and social sciences*, 2(6), 1251-1257.

23. Jamolov, F. N., Ravshanov, U. X., Atamurodov, B. N., Nabiyeu, M., Yanvarov, F., & Absalomov, S. (2024). GIDROUZELNI EKSPLOATATSIYA QILISH DAVRIDA XAVFSIZLIGI VA ISHONCHLILIGINI OSHIRISH (BESHARIQ GIDROUZELI MISOLIDA). *Лучшие интеллектуальные исследования*, 15(4), 124-129.

24. Фазлиев, Ж. Ш., Хаитова, И. И., Атамуродов, Б. Н., Рустамова, К. Б., & Шарипова, М. С. (2019). ТОМЧИЛАТИБ СУҒОРИШ ТЕХНОЛОГИЯСИНИ БОҒЛАРДА ЖОРИЙ ҚИЛИШНИНГ САМАРАДОРЛИГИ. *Интернаука*, (21-3), 78-79.

25. Juraev, A. K., Khamidov, M. K., Juraev, U. A., Atamurodov, B. N., Murodov, O. U., Rustamova, K. B., & Najmiddinov, M. M. (2024). The effect of deep softeners on irrigation, salt washing and cotton yield on soils with heavy mechanical composition. In *BIO Web of Conferences* (Vol. 103, p. 00017). EDP Sciences.

26. N., Atamurodov B., et al. "The Effectiveness of Farming in the Method of Hydroponics." *International Journal of Human Computing Studies*, vol. 3, no. 4, 2021, pp. 33-36, doi:10.31149/ijhcs.v3i4.2026.

27. Atamurodov, B. N., & Juraev, U. A. (2024). IMPACT OF INNOVATIVE SALINITY LEACHING AND IRRIGATION TECHNOLOGY ON SALT BALANCE IN EXPERIMENTAL FIELDS. *Академические исследования в современной науке*, 3(41), 151-153.